

Glassy trapping of manifolds in nonpotential random flows

Pierre Le Doussal¹ and Kay Jörg Wiese²

¹ CNRS-Laboratoire de Physique Théorique de l'Ecole Normale Supérieure, 24 rue Lhomond 75231 Paris*

² Fachbereich Physik, Universität GH Essen, 45117 Essen, Germany

We study the dynamics of polymers and elastic manifolds in non potential static random flows. We find that barriers are generated from combined effects of elasticity, disorder and thermal fluctuations. This leads to glassy trapping even in pure barrier-free divergenceless flows $v \stackrel{f \rightarrow 0}{\sim} f^\phi$ ($\phi > 1$). The physics is described by a new RG fixed point at finite temperature. We compute the anomalous roughness $R \sim L^\zeta$ and dynamical $t \sim L^z$ exponents for directed and isotropic manifolds.

Elastic structures in presence of quenched disorder exhibit glassy behaviour [1–3] such as pinning and very slow thermally activated motion of velocity $v \sim e^{-U(f)/T}$. Understanding these properties is crucial for numerous experimental systems, e.g. vortex lattices in superconductors, magnetic domains, solid friction and charge density waves [1,3]. For $D \leq 4$ the static system is usually in a disorder dominated glass state described by a $T = 0$ RG fixed point [4]. Barriers diverge at small force $U(f) \sim f^{-\mu}$ and large scales [1]. If $D \leq 2$ thermal fluctuations become important again. Periodic structures e.g. still exhibit a glass phase in $D = 2$, but with weaker logarithmic barriers $v \sim f^\phi$ described by $T > 0$ RG fixed points [5,6]. Up to now, the study of pinning and glassy dynamics has focused on *potential* systems, typically an elastic manifold in a random potential [7]. The existence of pinning and barriers in a *non potential* system was proposed recently in the context of vortex lattices driven over a disordered substrate [3]. Driven systems even when all microscopic forces are formally the gradient of a potential, are usually not so at a macroscopic level since energy is constantly pumped into the system. In [3] it was found that despite the non potential convective term present in the equation of motion, some glassy properties survive. Related observations were made in (mostly mean field) spin models [8,9]. Since little is known about general properties of non potential systems with disorder, it is of great interest to study other examples (and beyond mean field). An outstanding question is whether the effective temperature, which tends to be generated in these systems from the constant dissipation, does overwhelm the effect of static disorder.

In this Letter, we study a model of a polymer in a non potential static random flow (a “randomly driven polymer”) and its generalization to an elastic manifold. We work directly in finite dimension using dynamical RG. We study two cases: either the polymer is directed (each monomer sees a different flow) or it is isotropic (all monomers are in the same flow) see Fig. 1. Our main finding is that the physics is described by new RG fixed points with both $T > 0$ and finite disorder, leading to anomalous roughness ζ and glassy trapping by the flow

(with sublinear $v(f) \sim f^\phi$). Compared to the single particle studies [10], new features emerge such as the crucial role of internal elasticity in generating barriers. While being consistent with the Hartree results ($d \rightarrow \infty$) for long range (LR) disorder [9], our present study yields new universal fixed points for short range (SR) disorder. It also generalizes the dynamics of self avoiding manifolds [11] to a quenched disorder situation, though self avoidance is perturbatively less relevant in most regimes studied here. For real polymers in realistic flows ours is mostly a toy model which could be improved by including hydrodynamic forces. Nevertheless, some of the physics unveiled here will be present in more realistic models, such as the existence of preferred regions in the flows. Inhomogeneities in polymer distribution and non linearities in $v(f)$ could be investigated experimentally.




FIG. 1. elastic manifolds (polymers $D = 1$) in random flows: (a) directed polymer (b) isotropic chain.

Let us illustrate how elasticity (or entropy) leads to dynamical generation of barriers in a divergenceless flow. It is well known that for a single particle convected in such a flow, the stationary measure at $T > 0$ is spatially uniform and the drift velocity under a force simply $v = f$. Remarkably as soon as one considers two elastically coupled particles, preferred regions appear. Consider motion in the simple $2d$ flow ($v_x = -y$, $v_y = x$):

$$\begin{aligned} \dot{z}_1 &= c(z_2 - z_1) + i(\omega + \delta)z_1 + \eta_1 \\ \dot{z}_2 &= c(z_1 - z_2) + i(\omega - \delta)z_2 + \eta_2 \end{aligned} \quad (1)$$

$z_k = x_k + iy_k$ is the complex position of particle k , c the elastic coupling, $\langle \eta_i \eta_j^* \rangle = 4T\delta_{ij}$ is the thermal noise. Without disorder $\delta = 0$ the motion is just a collective rotation around the center (the matrix eigenvalues are $i\omega$ and $-2c + i\omega$). At $T > 0$ there is free collective diffusion and $V = f$ as before. As soon as $\delta > 0$ the zero mode disappears and the two particles converge towards the center (the eigenvalues are $\lambda_{\pm} = -c + i\omega \pm \sqrt{c^2 - \delta^2}$). This effect persists at $T > 0$ since one finds $\langle |z_k|^2 \rangle = 2T(c^2 + \delta^2)/(c\delta^2)$ ($k = 1, 2$) and thus there is a genuine bound state despite the divergenceless nature of the flow. The stationary conformation is twisted since $\langle z_1 z_2^* \rangle = 2T(c + i\delta)/\delta^2$. Extending (1) to a directed chain, one finds similar results (the decay towards the center at $T = 0$ is exactly the problem of spin depolarisation [12]). Thus a directed polymer in a flow will be attracted to regions where elastic energy and dissipation are smaller (though the precise balance remains to be understood). Presumably polymers in realistic flows will be repulsed by high vorticity regions. Non linear extensions of the above model will thus show the generation of barriers. Similar effects exist if all monomers see the same flow (isotropic chain). This is reminiscent of the effect discovered by Fauve and Thual [13]: mapping their Landau-Ginsburg function onto the complex position $z(x)$ of an elastic chain in a 2d non potential non linear flow, one sees that (even at $T = 0$) it may converge to stable localized (rotating and twisted) conformations $z(x, t) \rightarrow z_0(x)e^{i\omega t}$.

We now turn to our full disordered model where these effects can be studied quantitatively. We consider a manifold of internal dimension D parametrized by a d -component field $r^\alpha(x)$. The polymer corresponds to $D = 1$ (x labels the monomers), and a single particle to $D = 0$. We study the Langevin dynamics:

$$\eta \partial_t r_{xt}^\alpha = c \nabla_x^2 r_{xt}^\alpha + F_\alpha[r_{xt}, x] + \zeta_{xt}^\alpha \quad (2)$$

η is the friction, c the elastic coefficient and the gaussian thermal noise satisfies $\langle \zeta_{xt}^\alpha \zeta_{x't'}^\beta \rangle = 2\eta T \delta_{\alpha\beta} \delta(t - t') \delta^D(x - x')$. Angular brackets denote thermal averages and overbars disorder averages. $F_\alpha[r, x]$ is a gaussian quenched random force field of correlations:

$$\overline{F_\alpha[r, x] F_\beta[r', x']} = \Delta_{\alpha\beta}(r - r') h_{x-x'} \quad (3)$$

There are two main cases of interest. If the manifold is *directed* (e.g. a polymer oriented by an external field), then $h_{x-x'} = \delta^D(x - x')$. If the manifold is *isotropic*, (e.g. a gaussian chain in a static flow), the force field does not depend on the internal coordinate $F_\alpha[r, x] = F_\alpha[r]$ and $h_{x-x'} = 1$. We consider a statistically rotationally invariant force field with both a potential (L) (“electric”) and a divergence-free (T) part (“magnetic”). We thus study correlators $\Delta_{\alpha\beta}(r) = \int_K \Delta_K^{\alpha\beta} e^{iK \cdot r}$ with, at small K :

$$\Delta_K^{\alpha\beta} \sim K^{a-d} (g_L \frac{K_\alpha K_\beta}{K^2} + g_T (\delta_{\alpha\beta} - \frac{K_\alpha K_\beta}{K^2})) \quad (4)$$

with $\int_K = \int \frac{d^d K}{(2\pi)^d}$. We are mostly interested in *SR correlated forces* $a = d$ (force correlator decaying faster than r^{-d}). We also give results for *LR correlated forces* (force correlator decaying as r^{-a} with $a < d$). Note that *SR random potentials* $\Delta_{\alpha\beta}(K) \sim \tilde{g}_L K_\alpha K_\beta + \tilde{g}_T (\delta_{\alpha\beta} K^2 - K_\alpha K_\beta)$ (formally $a > d$) renormalize to the random force case, except in the potential case $g_T = 0$, much studied previously [1,2,4] which is preserved by RG.

We study the time translational invariant steady state. One defines the radius of gyration (or roughness) exponent $\langle (r_{xt} - r_{x't'})^2 \rangle \sim |x - x'|^{2\zeta}$, the single monomer diffusion exponent $\langle (r_{xt} - r_{x't'})^2 \rangle \sim |t - t'|^{2\nu}$ and assumes scaling behaviour $\langle (r_{xt} - r_{00})^2 \rangle \sim |x|^{2\zeta} \overline{B}(t/x^z)$ with $\zeta = z\nu$. The drift velocity under a small additional applied force f in (2) is $v \sim \langle r_{xt} \rangle / t$. We find $v \sim f^\phi$ at small f , with $\phi > 1$ and thus the polymer is trapped by the flow. Without disorder one has $\zeta_0 = (2 - D)/2$, $\nu_0 = (2 - D)/4$ for $D < 2$, and bounded fluctuations for $D > 2$.




FIG. 2. Regions in d, D plane where SR disorder is relevant for a directed manifold. Inset: Regions in a, d plane where SR or LR disorder is relevant for the directed polymer $D = 1$.

We start with simple Flory (i.e. dimensional) estimates. Balancing all terms in (2) (e.g. for directed manifolds it reads $r/t \sim r/x^2 \sim r^{-d/2} x^{-D/2}$) yields:

$$\zeta_F^{\text{dir}} = 2\nu_F^{\text{dir}} = \frac{4 - D}{2 + d} \quad , \quad \zeta_F^{\text{iso}} = 2\nu_F^{\text{iso}} = \frac{4}{2 + d} \quad (5)$$

for directed and isotropic manifolds, respectively (with $d \rightarrow a$ for LR disorder). This suggests that the manifold will be stretched by disorder [14]. We now study (2,3) using the dynamical generating functional $Z = \int Dr D\hat{r} e^{-S_0[r, \hat{r}] - S_{\text{int}}[r, \hat{r}] + if\hat{r}}$ with:

$$S_0 = \int d^D x dt \, i\hat{r}_{xt}^\alpha (\eta \partial_t - c \nabla_x^2) r_{xt}^\alpha - \eta T i\hat{r}_{xt}^\alpha i\hat{r}_{xt}^\alpha \quad (6)$$

$$S_{\text{int}} = -\frac{1}{2} \int_{x, x', t, t'} (i\hat{r}_{xt}^\alpha) (i\hat{r}_{x't'}^\beta) \Delta_{\alpha\beta}(r_{xt} - r_{x't'}) h_{x-x'}$$

We used dynamical RG methods, both via a Wilson RG scheme (presented here) as in [3,6] and via a field theoretical method using MOPE techniques as in [18,11] (detailed in [19]). The free propagator is $B(x, t) = \frac{1}{2D} \langle (r_{xt} - r_{00})^2 \rangle_0$ and the free response function is $DR(x, t) = \langle i\hat{r}_{00} \cdot r_{xt} \rangle_0$. (6, 4) is invariant under the rescaling $x = e^l x'$, $t = e^{z l} t'$, $r = e^{\zeta l} r'$, $\hat{r} = e^{(2-z-\zeta-D+\beta)l} \hat{r}'$, provided $\eta \rightarrow \eta e^{(2-z+\beta)l}$, $T \rightarrow T e^{(2-D-2\zeta+\beta)l}$, $c \rightarrow c e^{\beta l}$. Also $g_{T,L} \rightarrow g_{T,L} e^{(4-D+2\beta-(a+2)\zeta)l}$ (directed case) and $g_{T,L} \rightarrow g_{T,L} e^{(4+2\beta-(a+2)\zeta)l}$ (isotropic case) with $a = d$ everywhere for SR forces. There are generically *three* independent exponents z, ζ, β and from them one gets:

$$\nu = \zeta/z, \quad \phi = (z - \zeta)/(2 - \zeta + \beta) \quad (7)$$

using that $f \rightarrow f e^{(D+z+\alpha)l}$, $v \rightarrow v e^{(\zeta-z)l}$ under rescaling. Power counting at the Gaussian fixed point with no disorder ($z = 2, \beta = 0$ and $\zeta = (2-D)/2$), shows that disorder is relevant when (Fig.2) $d < d_c^{\text{dir}}$ for directed manifolds and $d < d_c^{\text{iso}}$ for isotropic manifolds (resp. $a < d_c$ for LR disorder), with:

$$d_c^{\text{dir}} = \frac{4}{2-D}, \quad d_c^{\text{iso}} = \frac{4+2D}{2-D} \quad (8)$$

Thus for directed polymers disorder is relevant for $d < 4$ and for isotropic polymers for $d < 6$. Power counting and symmetries show that the only relevant terms generated in perturbation theory are the one in (6). To lowest order the corrections to T, c and η read:

$$\begin{aligned} \delta T &= \frac{(d-1)g_T}{\eta d} \int_{\tau>0, y, K} h_y K^{a-d} (e^{-K^2 B(y, \tau)} - e^{-K^2 B_\infty}) \\ \delta c &= \frac{g_L}{2dD} \int_{y, K} y^2 h_y K^{a-d} (e^{-K^2 B(y, 0)} - e^{-K^2 B_\infty}) \end{aligned} \quad (9)$$

and $(d-1)\delta\eta = (g_L/g_T)\eta\delta T$ (where $B_\infty = \int_k \frac{T}{ck^2}$ is IR infinite for $d \leq d_c$) [20]. To simplify the above expressions we have used that the free propagator satisfies the FDT relation $\theta(t) \frac{d}{dt} B(x, t) = TR(x, t)$. We note that a useful feature of the present model is that although the FDT relation is not obeyed by the exact Green's functions (since disorder is non potential) it is within each scale of the RG. The FDT violation will simply appear as a non trivial renormalization of T . The relevant corrections to disorder read:

$$\begin{aligned} \delta\Delta_P^{\alpha\beta} &= \int_{x, y, \tau, \tau'} \int_{K, K'=P-K} (\Delta_K^{\alpha\beta} (K \cdot \Delta_{K'} \cdot K) \\ &+ (K' \cdot \Delta_K^\alpha) (K \cdot \Delta_{K'}^\beta)) R_{x, \tau} R_{y, \tau'} h_{x-y} e^{K \cdot K' (B_{x, \tau} + B_{y, \tau'})} \end{aligned} \quad (10)$$

The small P expansion in (10) is well behaved and because of analyticity the only divergent terms generated are proportional to $\delta_{\alpha\beta}$ yielding $\delta g_L = \delta g_T$. One can thus take the limit $P \rightarrow 0$ ($K' = -K$ in (10)). One checks explicitly from (10) that (i) starting with $\Delta_{\alpha\beta}(K) \sim \tilde{g}_L K_\alpha K_\beta + \tilde{g}_T (\delta_{\alpha\beta} K^2 - K_\alpha K_\beta)$ with both

$\tilde{g}_T > 0$ and $\tilde{g}_L > 0$ one generates SR random forces and that (ii) a divergence free random force field does not remain so. The final RG equations have the form (specified to various cases below and in Table 1):

$$\frac{dc}{cdl} = \beta + A_c \bar{g}_L, \quad \frac{d\eta}{\eta dl} = 2 - z + \beta + A_\eta \bar{g}_L \quad (11)$$

$$\frac{dT}{T dl} = 2 - D - 2\zeta + \beta + A_T \bar{g}_T \quad (12)$$

$$\frac{d\bar{g}_L}{dl} = \epsilon \bar{g}_L - (B - A) \bar{g}_L \bar{g}_T + E \bar{g}_L^2 \quad (13)$$

$$\frac{d\bar{g}_T}{dl} = \epsilon \bar{g}_T + (A + E) \bar{g}_L \bar{g}_T - B \bar{g}_T^2 \quad (14)$$




FIG. 3. Examples of RG flow diagrams. (a) directed manifolds with SR disorder. The physics is controlled by the fixed point I^* at $g_T = g_L$. (b) isotropic manifolds LR disorder. The flow is along straight lines. There is a line of fixed points L and an apparent separatrix. $\zeta - \zeta_0$ changes sign upon increasing g_L/g_T , suggesting progressive localization.

(i) *Directed manifolds, SR disorder*: $\epsilon = 2 - (2-D)\frac{d}{2}$. Then $h_y = \delta^D(y)$ and there is no correction to the elastic coefficient $\delta c = 0$ (guaranteed by the statistical tilt symmetry [3] for SR and LR disorder). Thus $\beta = 0$ and there are only two independent exponents. The dimensionless coupling [17] constant is $\bar{g}_{T,L} = C \Lambda^\epsilon g_{T,L} / T^{(d+2)/2}$. The RG flow is depicted in Fig.3. There is a globally attractive isotropic fixed point I^* at $\bar{g}_L = \bar{g}_T = \epsilon/(B-A)$ (note that $B > A$) and the line $\bar{g}_L = \bar{g}_T$ is preserved. The potential line $g_T = 0$ is preserved with a flow to strong coupling. There is another *apparent* fixed point at $g_L = 0, g_T = \epsilon/B$, attractive along the line $g_L = 0$ (divergenceless flows). However for $D > 0$ this line is not in the physical domain since, as discussed above, a finite g_L term is generated and one flows to I^* . At I^* one gets

$$z = 2 + \epsilon/a_{D^*}, \quad \zeta = \frac{2-D}{2} + \epsilon/(2b_{D^*}) \quad (15)$$

$$b_D = \frac{4-D}{2-D} - \frac{4}{D} \left(\frac{1}{2\Gamma[D/2]} \right)^{2/(2-D)}, \quad a_D = \frac{2+D}{2-D} b_D$$

and ν and ϕ using (7). The expansion can be carried from any point of the line $d^* = 4/(2-D^*)$. Optimizing over D^* as in [18] yields the estimates $\zeta = 0.625 \pm 0.02$, $z = 2.085 \pm 0.02$, $\phi = 1.061 \pm 0.015$ ($d=3$) and $\zeta = 0.9 \pm 0.1$,

$z = 2.2 \pm 0.1$, $\phi = 1.182 \pm 0.08$ ($d=2$). For small ϵ , $\zeta > \zeta_F$ and Flory is thus likely to be a lower bound [23]

(ii) *directed manifolds, LR disorder*: $\epsilon = 2 - \frac{(2-D)a}{2}$ with $a < d$. The novelty is $\delta g_{L,R} = 0$ since the LR part of the disorder is not renormalized [15]. The dimensionless coupling [17] $\bar{g}_{L,T} = C g_{L,T} / T^{(a+2)/2}$ will experience non trivial renormalization only because T renormalizes. Thus the ratio $\frac{\bar{g}_L(l)}{\bar{g}_T(l)} = \frac{g_L}{g_T}$ is preserved which leads to a *line of fixed points*. Thus $\bar{g}_T(l) \rightarrow g_T^* = \epsilon/B$ and $\bar{g}_L(l) \rightarrow \frac{g_L}{g_T} g_T^*$. This yields the continuously varying exponent $z = 2 + 2 \frac{g_L}{g_T} \frac{\epsilon}{(2+a)(d-1)}$ and $\zeta = \zeta_F$ holds *to all orders* in ϵ in the LR case since neither the vertex nor c renormalize. As conjectured in [9] ν is found identical to the Hartree approximation (formula (12) [22] of [9]).

(iii) *isotropic manifolds (SR and LR disorder)*: Then $\epsilon = D + 2 - a \frac{2-D}{2}$ ($a = d$ for SR disorder). The novelty is a renormalization of the elasticity of the manifold (it becomes stiffer $\delta c > 0$) and thus a third non trivial exponent β . The dimensionless constant is $\bar{g}_{T,L} = C \frac{g_{T,L}}{T^{\frac{2+a}{2} c^{\frac{2-a}{2}}}}$. For SR disorder the RG flow is similar to the directed case. There is a fully attractive (isotropic disorder) fixed point I^* at $\bar{g}_T = \bar{g}_L = 1/(B - E - A)$. As before, LR disorder is not renormalized, the flow lines are $\frac{\bar{g}_L(l)}{\bar{g}_T(l)} = \frac{g_L}{g_T} = k$ and there is a line of fixed points $g_T = \epsilon/(B - kE)$, $g_L = k g_T$ parametrized by k . The general formula for the exponent is $\zeta = (2 - D)/2 + \epsilon(A_T - kA_c)/2(B - kE - A)$, $z = 2 + k\epsilon(A_\eta - A_c)/(B - kE - A)$ and $\beta = -k\epsilon A_c/(B - kE - A)$, with the values given in Table 1 (and set $k = 1$ for SR). For SR disorder this yields for $D = 1$, $\zeta = 0.5 + 0.13\epsilon$, $z = 2 + 0.04\epsilon$, $\beta = -0.015\epsilon$. Extrapolations as in [18] suggest $\zeta = 0.85 \pm 0.1$, $z = 2.1 \pm 0.1$, $\beta = -0.025 \pm 0.01$ for polymers in $d = 3$. Note that for LR disorder the line of fixed points is not parallel to the axis any more (see Fig. 3), and thus there is *no fixed point* for $k > k_c = B/E$. There is even a value $k^* = \frac{a-2}{a+2} k_c < k_c$ where $\zeta - \zeta_0$ changes sign. For $k^* < k < k_c$ the localization effects dominate. Together with the apparent separatrix in the flow, this suggests two phases (a localized one and another one with a continuously varying ζ). It would be interesting to check this scenario by non perturbative methods (such as in [9]).

Thus polymers (and manifolds) in static flows are described by non trivial RG fixed points at finite disorder. The main difference with conventional static (potential) glasses is that a temperature is generated which weakens disorder (and also renders the fixed point perturbatively accessible - the coupling constant being $g/T^{(d+2)/2}$!). The main similarity lies in the sublinear $v(f)$ response reminiscent of the marginal glass in the statics [5]. Numerical simulations would be helpful to check the present results and further investigate the physics of this phase [24]. The competition between localization and driving found in some cases for isotropic manifolds (i.e correlated disorder) also deserves further studies.

We thank L. Cugliandolo, J. Kurchan, T. Giamarchi and L. Schäfer for discussions.

-
- * Laboratoire Propre du CNRS, associé à l'ENS et à l'Université Paris-Sud.
- [1] G. Blatter *et al.*, Rev. Mod. Phys. **66**, 1125 (1994).
 - [2] M. Kardar cond-mat/9704172 and cond-mat/9507019.
 - [3] T. Giamarchi, P. Le Doussal, Phys. Rev. Lett. **76** 3408 (1996), cond-mat/9705096, cond-mat/9708085.
 - [4] D. S. Fisher, Phys. Rev. Lett. **56**, 1964 (1986).
 - [5] J. L. Cardy and S. Ostlund, Phys. Rev. B **25**, 6899 (1982),
 - [6] D. Carpentier and P. Le Doussal, Phys. Rev. B **55**, 12128 (1997) and references therein.
 - [7] or hamiltonian systems (which have similar properties).
 - [8] L. F. Cugliandolo, J. Kurchan, P. Le Doussal, and L. Peliti, Phys. Rev. Lett. **78**, 350 (1997).
 - [9] P. Le Doussal, L. F. Cugliandolo, L. Peliti Europhys. Lett. **39** 111 (1997).
 - [10] J.P. Bouchaud, A. Comtet, A. Georges, P. Le Doussal, J. Physique (Paris) **48** 1445 (1987), **49** 369 (1988), J. Honkonen and E. Karjalainen, J. Phys. A **21**, 4217 (1988) and references therein.
 - [11] K. J. Wiese, cond-mat/9702020 and cond-mat/9702023
 - [12] P.P. Mitra, P. Le Doussal, Phys. Rev. B **44** 12035 (1991).
 - [13] O. Thual and S. Fauve J. Phys. France **49** 1829 (1988).
 - [14] however $\zeta = \zeta_F$ is exact *only* for directed manifolds with LR correlations with also $\nu = \nu_F$ $g_L = 0$ and $d \rightarrow \infty$ (while for $D = 0$ and $g_L = 0$, $\nu = \nu_F$ is *always* exact [9]).
 - [15] (10) renormalizes only the SR part of disorder and becomes important only in the region of the (a, d) plane where the crossover [10] LR-SR occurs (not studied here).
 - [16] We have extended the calculation of [9] and obtained the ϕ exponent for Hartree in agreement with (7).
 - [17] We use $C = \frac{\Gamma[a/2]}{\Gamma[d/2]} (4\pi)^{\frac{a-d}{2}-1} (\frac{2-D}{2})^{a/2}$ and $C' = (2 - D)^{a/2} \frac{\Gamma[a/2]}{\Gamma[d/2]} S_D^{1+\frac{a}{2}} (4\pi)^{-d/2}$ with $a = d$ for SR disorder.
 - [18] F. David and K.J. Wiese, Phys. Rev. Lett. **76** 4564 (1996), K.J. Wiese and F. David, Nucl. Phys. B **487** 529 (1997).
 - [19] K. Wiese, P. Le Doussal, to be published
 - [20] (9,10) are computed using $B(x, 0) = T|x|^{2-D}/((2 - D)S_D c)$, $B(0, t) = 2T|\mu t|^{\frac{2-D}{2}}/((2 - D)(4\pi c)^{D/2})$ with $S_D = 2\pi^{D/2}/\Gamma(D/2)$ and $\int_K e^{-K^2 B} K^{a-d} = (4\pi)^{-d/2} \frac{\Gamma[a/2]}{\Gamma[d/2]} B^{-a/2}$.
 - [21] Continuing (15) to $D = 0$ yields $\zeta = 1 + \frac{\epsilon}{2}$, $z = 2 + \epsilon$ (with $\epsilon = 2 - d$). This yields $\nu = \zeta/z = 1/2 + O(\epsilon^2)$ and $\phi = 1 + \epsilon$ at I^* which is the correct known result for the particle, obtained via a different RG [10].
 - [22] setting $g_2 \rightarrow g_L/d$, $g_1 - g_2 \rightarrow (1 - \frac{1}{d})g_T$ and $\epsilon \rightarrow \epsilon/(2 - D)$.
 - [23] Elasticity slows down diffusion compared to a single particle since e.g. $\nu(D = 1) < \nu(D = 0)$.
 - [24] A scenario to be checked is whether the hierarchy of wells within wells of a static glass is interrupted here at some

random level by the nonpotential forces (a mechanism to generate wells with algebraic distribution of release time)

and if there are Levy walk type of jumps between trapped configurations.

	A_c	A_η	A_T	B	E	A
dirSR	0	$\frac{2}{d}$	$\frac{2(d-1)}{d}$	$\frac{(2+d)}{2} A_T$	0	$\frac{16\pi(d-1)}{d(d-2)(2-D)} \left(\frac{S_D}{4\pi}\right)^{d/2}$
dirLR	0	$\frac{2}{d}$	$\frac{2(d-1)}{d}$	$\frac{(2+a)}{2} A_T$	0	0
isoSR	$\frac{1}{2dD}$	$\frac{I_d}{d}$	$\frac{(d-1)I_d}{d}$	$\frac{d+2}{2} A_T$	$\frac{d-2}{2} A_c$	$\frac{2(d-1)\Gamma[\frac{D}{2-D}]^2}{d(d-2)(2-D)^2\Gamma[\frac{2D}{2-D}]}$
isoLR	$\frac{1}{2dD}$	$\frac{I_a}{d}$	$\frac{(d-1)I_a}{d}$	$\frac{a+2}{2} A_T$	$\frac{a-2}{2} A_c$	0

Table I: Coefficients of the RG equation (11) for cases studied here (one must set $d = d^*$ and $D = D^*$). One denotes $I_a = \int_0^{+\infty} dt \tilde{B}(1, t)^{-a/2}$ with $\tilde{B}(1, t) = \frac{1}{\Gamma[D/2]}((4t)^{(2-D)/2} e^{-1/(4t)} + \Gamma[D/2, 1/(4t)])$, with $\tilde{I}_{d=6} \approx 1.7935$,